# **COVID-19 VACCINE BOOSTER INFO**

moderna

#### **WHO**

- 65 or older;
- 18 or older with an underlying medical condition or living in a long-term care setting;
- 18 or older working in a high-risk setting

#### **WHEN**

• At least 6 months after initial series

## DOSE

• One shot with the same dosage as the original series (30 micrograms)

## **MIX & MATCH**

· Yes, at least 6 months after receiving initial series

## STORAGE REQUIREMENTS

- Ultra-cold Freezer: between -90 degrees Celsius and -60 degrees Celsius until expiration date
- Freezer: between -25 degrees Celsius and -15 degrees Celsius for up to 2 weeks
- Refrigerator: between 2 degrees Celsius and 8 degrees Celsius for up to 31 days. Contact manufacturer after 31 days

## **COMMON SIDE FEFECTS**

 Pain at injection site, fatigue, muscle pain, headache, chills, swollen lymph nodes in the same arm as the injection site, fever, nausea and vomiting

# **WHO**

- 65 or older;
- 18 or older with an underlying medical condition or living in a long-term care setting;
- 18 or older working in a high-risk setting

## **WHEN**

· At least 6 months after initial series

## DOSE

• One shot with half the dosage of the original series (50 micrograms)

#### **MIX & MATCH**

· Yes, at least 6 months after receiving initial series

## STORAGE REQUIREMENTS

- Freezer: between -50 degrees Celsius and -15 degrees Celsius until expiration date
- Refrigerator: between 2 degrees Celsius and 8 degrees Celsius for up to 30 days. Contact the manufacturer after 30 days

## **COMMON SIDE EFFECTS**

• Pain at injection site, fatigue, muscle pain, headache, fever, chills and nausea

# **WHO**

· Anyone 18 and older that initially received a J&J shot

# WHEN

At least 2 months after initial shot

Johnson Johnson

# DOSE

• One shot with the same dosage as the original shot (0.5 mL)

## **MIX & MATCH**

· Yes, at least 2 months after receiving initial shot

## STORAGE REQUIREMENTS

• Refrigerator: between 2 degrees Celsius and 8 degrees Celsius until expiration date

## **COMMON SIDE EFFECTS**

· Pain at injection site, fatigue, muscle pain, headache, fever, chills and nausea

